

POST-SURGERY INSTRUCTIONS: BRACHIOPLASTY

Once your surgery is completed, you must follow all the instructions given to you in order to heal properly and have a good outcome.

The following instructions are your obligation. Use this as a checklist of progress as you heal. Included are normal post-surgical experiences and key health considerations that may be a cause of concern.

TYPICAL POST-OPERATIVE SYMPTOMS

Typical symptoms of Brachioplasty and signs to watch for following arm lift surgery include the following:

Tightness and stiffness in the arm and underarm region: Bruising, swelling and redness: Tingling, burning or intermittent shooting pain: These are normal experiences as the skin, tissues and sensory nerves heal. Pain medication and muscle relaxants will help you cope with any discomfort. If you have drains, you may experience additional localized discomfort. **Consistent sharp pain should be reported to our office immediately.**

Skin firmness, hypersensitivity or lack of sensitivity: This is normal and will gradually resolve over time.

Shiny skin or any itchy feeling: Swelling can cause the skin of your arms to appear shiny. As the healing process advances, you may also find a mild to severe itchy feeling in your upper arms. An antihistamine like Benadryl can help to alleviate severe, constant itchiness. **If the skin becomes red and hot to the touch, contact our office immediately.**

Asymmetry, the arms look different, or heal differently: Your arms may look or feel quite different from one another in the days following surgery. This is normal; no two arms in nature are perfectly symmetrical or following surgery.

CALL THE OFFICE IMMEDIATELY IF YOU EXPERIENCE ANY OF THE FOLLOWING:

- **A high fever, (over 101°) severe nausea and vomiting, continued dizziness or incoherent behavior, such as hallucinations.**
- **Any pain that cannot be controlled by your pain medication.**
- **Bright red skin that is hot to the touch.**
- **Excessive bleeding or fluid seeping through the incisions.**
- **A severely misshapen upper arm or bruising that is localized to one specific point of the arm.**

To alleviate any discomfort, and to reduce swelling, you may apply cool, not cold compresses to the treated region. Crushed ice or ice packs must be wrapped in a towel before being applied to the skin. Do not apply ice or anything frozen directly to the skin. Apply cool compresses, for no longer than 20-minute intervals.

DAY OF SURGERY INSTRUCTIONS

You will only be released to the care of a responsible adult. All of these instructions must be clear to the adult who will monitor your health and support you, around the clock during the first 24 hours following surgery.

Rest, but not bed rest. While rest is important in the early stages of healing, equally important is that you are ambulatory, meaning that you are walking under your own strength. Spend 10 minutes every 2 hours engaged in light walking indoors as you recover.

Recline, do not lie down flat. This will be more comfortable for you, and can reduce swelling. Always keep your head elevated. Do not bend forward or over. Elevate your arms on pillows.

Good nutrition. Fluids are critical following surgery. Stick to non-carbonated, non-alcoholic, caffeine-free and green tea-free beverages including fruit juices and water, milk and yogurt drinks. You must consume at least 8 ounces of fluid every 2 hours. Stick with soft, bland, nutritious food for the first 24 hours.

DO NOT Change your incision dressings. Your incisions will seep fluid and some blood for a short time after surgery. Keep dressings clean and dry. If dressings become saturated, remove gauze and replace. **Replace any compressions garments over the gauze.** If you have a drain placed in your incisions, carefully follow the instructions for drain care and record drained fluid on the **Drain Care Instructions and Log.**

Wear your compression garment or elastic wraps around the clock. Follow the instructions specifically, and only remove any compression to cleanse your incision or to empty any drains.

DO NOT SMOKE. Smoking can greatly impair your safety prior to surgery and your ability to heal following surgery. You must not smoke.

Relax. Do not engage in any stressful activities. Do not lift, push or pull anything. Take care of no one, and let others tend to you.

TWO TO SEVEN DAYS FOLLOWING SURGERY

During this time you will progress as each day passes. Ease into your daily activities. You will receive clearance to

begin driving or return to work at your post-operative visit.

- **Your dressings will be removed in the office 1-2 days after surgery**
- **Continue to cleanse wounds as directed; you may shower after the dressings are removed.** Make sure a responsible adult is home with you when you take a shower as you may become light-headed. Take a warm, not hot shower. Do not take a bath. Limit your shower to 10 minutes. A cotton swab with peroxide is appropriate for cleansing incisions. Do not remove any steri-strips over your stitches. Apply anti-bacterial ointment over the steri-strips, and then apply 4x4 gauze pads over your incisions. Do not remove any steri-strips. Do not rub your incisions. You may apply a fragrance free moisturizer to the arm and surrounding skin, however not on your incisions.
- **Take antibiotic medications and supplements as directed.** Take pain medication and muscle relaxants only as needed. You may wish to switch from prescriptive pain medication to acetaminophen or ibuprofen.
- **Continue to wear your elastic or compression wraps around the clock.**
- **Do not resume any exercise other than regular walking.** Walking is essential every day to prevent the formation of blood clots.
- **No sun exposure.** If you plan to go outdoors for any reason, wear long sleeves. Avoid any direct sun exposure.
- **Maintain a healthy diet. Do not smoke. Do not consume alcohol.**

ONE TO FOUR WEEKS FOLLOWING SURGERY

As you resume your normal daily activities, you must continue proper care and healing.

- **Continue your wound care as directed.**
- **Refrain from weight-bearing exercise, or lifting anything over your head.** You may begin range of motion exercises but not with any weight, pressure or resistance of any kind.
- **Continue to wear your elastic or compression wrap as directed.**
- **Practice good sun protection.** Do not expose your arms to direct sunlight. If you are outdoors, apply at least an SPF 30 to the arms at least 30 minutes prior to sun exposure and wear long sleeves. The skin of your arms is highly susceptible to sunburn or the formation of irregular, darkened pigmentation.

SIX WEEKS FOLLOWING SURGERY

Healing will progress and your arms will settle into a more final shape and position.

- **You may ease into your regular fitness routine.** However realize that your upper body may require some time to return to previous strength.
- **Discomfort or tightness and tingling of the skin and upper arm will resolve.**
- **No need to resume smoking.** You have now gone 8 weeks (2 weeks prior to surgery and 6 weeks following) without a cigarette. For your long-term health, there is no need to resume smoking.

YOUR FIRST YEAR

- **Continue healthy nutrition, fitness and sun protection.**
- **Your scars will continue to refine.** If they become raised, red or thickened, or appear to widen, contact our office. Early intervention is important to achieving well-healed scars. Scars are generally refined to fine incision lines one year after surgery.
- **A one-year post surgery follow-up is recommended.** However you may call our office at any time with your concerns or for needed follow-up.

Your body will change with age. The appearance of your arms will change too. Although the outcomes of Brachioplasty are generally permanent, any significant weight gain or loss, as well as the normal influences of aging can cause changes to your appearance. You may wish to undergo revision surgery at a later date to help maintain your appearance throughout life. Contact our office with any of your questions or concerns, at any time.

I have read and understand all of the above instructions. I understand that following these instructions is solely my responsibility. I understand that it is also my responsibility to ask Dr. Castor and/or his staff any questions I have related to these instructions or about my procedure, health and healing.